

A NOTE ON ELLIPTIC PLANE CURVES WITH FIXED j -INVARIANT

RAHUL PANDHARIPANDE

10 May 1995

1. SUMMARY

Let N_d be the number of irreducible, reduced, nodal, degree d *rational plane curves* passing through $3d - 1$ general points in the complex projective plane $\mathbf{P}^2$. The numbers N_d satisfy a beautiful recursion relation ([K-M], [R-T]):

$$N_1 = 1$$

$$\forall d > 1, \quad N_d = \sum_{i+j=d, i,j>0} N_i N_j \left(i^2 j^2 \binom{3d-4}{3i-2} - i^3 j \binom{3d-4}{3i-1} \right).$$

Let $E_{d,j}$ be the number of irreducible, reduced, nodal, degree d , *elliptic plane curves* with fixed j -invariant j passing through $3d - 1$ general points $\mathbf{P}^2$. $E_{d,j}$ is defined for $d \geq 3$ and $\infty \neq j \in \overline{M}_{1,1}$. In this note, the following relations are established:

$$\begin{aligned} \forall j \neq 0, 1728, \infty, \quad E_{d,j} &= \binom{d-1}{2} N_d, \\ j = 0, \quad E_{d,0} &= \frac{1}{3} \binom{d-1}{2} N_d, \\ j = 1728, \quad E_{d,1728} &= \frac{1}{2} \binom{d-1}{2} N_d. \end{aligned}$$

If $d \equiv 0 \pmod{3}$, then $3 \nmid \binom{d-1}{2}$. Since $E_{3\hat{d},0}$ is an integer, $N_{3\hat{d}} \equiv 0 \pmod{3}$ for $\hat{d} \geq 1$. In fact, a check of values in [D-I] shows $N_d \equiv 0 \pmod{3}$ if and only if $d \equiv 0 \pmod{3}$ for $3 \leq d \leq 12$. P. Aluffi has calculated $E_{3,j}$ for $j < \infty$ in [A]. Aluffi's results agree with the above formulas.

Thanks are due to Y. Ruan for discussions on Gromov-Witten invariants and quantum cohomology. The question of determining the numbers $E_{d,j}$ was first considered by the author in a conversation with Y. Ruan.

2. KONTSEVICH'S SPACE OF STABLE MAPS

2.1. The Quasi-Projective Subvarieties $U_C(\Gamma, c, \overline{w})$, $U_{j=\infty}(\Lambda, \overline{w})$. Fix $d \geq 3$ for the entire paper. Let C be a nonsingular elliptic curve or an irreducible, nodal rational curve of arithmetic genus 1. Consider the coarse moduli space of $3d - 1$ -pointed stable maps from C to $\mathbf{P}^2$ of degree $d \geq 3$, $\overline{M}_{C, 3d-1}(\mathbf{P}^2, d)$. For convenience, the notation $\overline{M}_C(d) = \overline{M}_{C, 3d-1}(\mathbf{P}^2, d)$ will be used. Let $S_d = \{1, 2, \dots, 3d - 1\}$ be the marking set. Constructions of $\overline{M}_C(d)$ can be found in [Al], [K], [P]. Let Γ be a tree consisting of a distinguished vertex c , $k \geq 0$ other vertices $v_1, \dots, v_k$, and $3d - 1$ marked legs. Let $0 \leq e \leq d$. Weight the vertex c by e . Let $w_1, \dots, w_k$ be non-negative integral weights of the vertices $v_1, \dots, v_k$ satisfying

$$e + w_1 + \dots + w_k = d.$$

Denote the weighting by $\overline{w} = (e, w_1, \dots, w_k)$. The marked, weighted tree with distinguished vertex $(\Gamma, c, \overline{w})$ is stable if the following implication holds for all $1 \leq i \leq k$:

$$w_i = 0 \Rightarrow \text{valence}(v_i) \geq 3.$$

Two marked, weighted trees with distinguished vertex $(\Gamma, c, \overline{w})$ and $(\Gamma', c', \overline{w}')$ are isomorphic if there is an isomorphism of marked trees $\Gamma \rightarrow \Gamma'$ sending c to c' and respecting the weights.

A quasi-projective subvariety $U_C(\Gamma, c, \overline{w})$ of $\overline{M}_C(d)$ is associated to each isomorphism class of stable, marked, weighted graph with distinguished vertex $(\Gamma, c, \overline{w})$. $U_C(\Gamma, c, \overline{w})$ consists of stable maps $\mu : (D, p_1, \dots, p_{3d-1}) \rightarrow \mathbf{P}^2$ satisfying the following conditions. The domain D is equal to a union:

$$D = C \cup \mathbf{P}_1^1 \cup \dots \cup \mathbf{P}_k^1.$$

The marked, weighted dual graph with distinguished vertex of the map μ is isomorphic to $(\Gamma, c, \overline{w})$. The distinguished vertex of the dual graph of μ corresponds to the (unique) component of D isomorphic to C . Weights of the dual graph of μ are obtained by the degree of μ on the components. Note $U_C(\Gamma, c, \overline{w}) = \emptyset$ if and only if $e = 1$.

Let $(\Gamma, c, \overline{w})$ be a stable, marked, weighted tree with distinguished vertex. Assume $e \neq 1$. The dimension of $U_C(\Gamma, c, \overline{w})$ is determined as follows. If $e \geq 2$, then

$$\dim U_C(\Gamma, c, \overline{w}) = 6d - 2 - k.$$

If $e = 0$, then

$$\dim U_C(\Gamma, c, \overline{w}) = 6d - k$$

(where k is the number of non-distinguished vertices of Γ). These calculations are straightforward.

Let C be a nonsingular elliptic curve. Every stable map in $\overline{M}_C(d)$ has domain obtained by attaching a finite number of marked trees to C . By the definition of tree and map stability:

$$\bigcup_{(\Gamma, c, \overline{w})} U_C(\Gamma, c, \overline{w}) = \overline{M}_C(d).$$

Let C be an irreducible, 1-nodal rational curve. The quasi-projective varieties $U_C(\Gamma, c, \overline{w})$ do not cover $\overline{M}_C(d)$. The curve C can degenerate into a simple circuit of $\mathbf{P}^1$'s. Let Λ be a graph with 1 circuit (1^{st} Betti number equal to 1, no self edges), $k \geq 1$ vertices $v_0, \dots, v_k$, and $3d - 1$ marked legs. Note the different vertex numbering convention. At least 2 vertices are required to make a circuit, so $k \geq 1$. Let $w_0, w_1, \dots, w_k$ be non-negative, integral weights summing to d . The marked, weighted graph with 1 circuit $(\Lambda, \overline{w})$ is stable if each zero weighted vertex has valence at least 3. A quasi-projective subvariety $U_C(\Lambda, \overline{w})$ of $\overline{M}_C(d)$ is associated to each isomorphism class of stable, marked, weighted graph with 1 circuit $(\Lambda, \overline{w})$. $U_C(\Lambda, \overline{w})$ consists of stable maps with marked, weighted dual graphs isomorphic to $(\Lambda, \overline{w})$. The union

$$\bigcup_{(\Gamma, c, \overline{w})} U_C(\Gamma, c, \overline{w}) \cup \bigcup_{(\Lambda, \overline{w})} U_C(\Lambda, \overline{w}) = \overline{M}_C(d)$$

holds by the definition of stability.

Finally, the dimensions of the loci $U_C(\Lambda, \overline{w})$ will be required. Let $(\Lambda, \overline{w})$ be a stable, marked, weighted graph with 1 circuit. Let $c_1, \dots, c_l$ be the unique circuit of vertices of Λ . Let e be the sum of the weights of the circuit vertices. $U_C(\Lambda, \overline{w}) = \emptyset$ if and only if $e = 1$. If $e \geq 2$, then

$$\dim U_C(\Lambda, \overline{w}) = 6d - 2 - k.$$

If $e = 0$, then

$$\dim U_C(\Lambda, \overline{w}) = 6d - k$$

(where $k + 1$ is the total number of vertices of Λ). Again, these results are straightforward.

2.2. The Component $\overline{W}_1(d)$. Let $\overline{M}_1(d) = \overline{M}_{1, 3d-1}(\mathbf{P}^2, d)$ be Kontsevich's space of $3d - 1$ -pointed stable maps from genus 1 curves to $\mathbf{P}^2$. There is canonical morphism

$$\pi : \overline{M}_1(d) \rightarrow \overline{M}_{1,1}$$

obtained by forgetting the map and all the markings except $1 \in S_d$ (the $3d - 1$ possible choices of marking in S_d all yield the same morphism π). Let $j \in \overline{M}_{1,1}$. By the universal properties of the moduli spaces, there is a canonical bijection

$$\overline{M}_{C_j}(d) \rightarrow \pi^{-1}(j)$$

where C_j is the elliptic curve (possibly nodal rational) with j -invariant j . When $j \in \overline{M}_{1,1}$ is automorphism-free, this bijection is an isomorphism. For $j = 0, 1728$, the scheme theoretic fiber $\pi^{-1}(j)$ is nonreduced. Define an open locus $W_1(d) \subset \overline{M}_1(d)$ by $[\mu : (D, p_1, \dots, p_{3d-1}) \rightarrow \mathbf{P}^2] \in W_1(d)$ if and only if D is irreducible. By considering the natural tautological spaces over the universal Picard variety of degree d line bundles over $M_{1,3d-1}$, it is easily seen that $W_1(d)$ is a reduced, irreducible open set of dimension $6d - 1$. Let $\overline{W}_1(d)$ be the closure of $W_1(d)$ in $\overline{M}_1(d)$.

3. A DEFORMATION RESULT

Let Φ be a stable, marked, weighted tree with distinguished vertex c determined by the data: $k = 1$, $(e, w_1) = (0, d)$. There are 2^{3d-1} isomorphism classes of such Φ determined by the marking distribution. Let $j \in \overline{M}_{1,1}$. The dimension of $U_j(\Phi)$ is $6d - 1$. A point $[\mu] \in U_j(\Phi)$ has domain $C_j \cup \mathbf{P}^1$. There are $3d - 1$ dimensions of the map $\mu|_{\mathbf{P}^1} : \mathbf{P}^1 \rightarrow \mathbf{P}^2$. The incidence point $p = C_j \cap \mathbf{P}^1$ moves in a 1-dimensional family on $\mathbf{P}^1$. The remaining $3d - 1$ markings move in $3d - 1$ dimensions on C_j and $\mathbf{P}^1$ (specified by the marking distribution). $6d - 1 = 3d - 1 + 1 + 3d - 1$. A technical result is needed in the computation of the numbers $E_{d,j}$.

Lemma 1. *Let $I(\Phi, j) = \overline{W}_1(d) \cap U_j(\Phi) \subset \overline{M}_1(d)$. The dimension of $I(\Phi, j)$ is bounded by $\dim I(\Phi, j) \leq 6d - 3$.*

Proof. Let $[\mu] \in I(\Phi, j)$ be a point. Let $D = C_j \cup \mathbf{P}^1$ be the domain of μ as above. The following condition will be shown to hold: the linear series on $\mathbf{P}^1$ determined by $\mu|_{\mathbf{P}^1}$ has vanishing sequence $\{0, \geq 2, *\}$ at the incidence point $p = C_j \cap \mathbf{P}^1$. The existence of a point with vanishing sequence $\{0, \geq 2, *\}$ is a 1-dimensional condition on the linear series. The condition that the incidence point p has this vanishing sequence is an additional 1-dimensional constraint on p . Therefore, the dimension of $I(\Phi, j)$ is at most $6d - 1 - 1 - 1 = 6d - 3$. The vanishing sequence $\{0, \geq 2, *\}$ is equivalent to $d(\mu|_{\mathbf{P}^1}) = 0$ at p .

It remains to establish the vanishing sequence $\{0, \geq 2, *\}$ at p . This result is easily seen in explicit holomorphic coordinates. Let Δ_t be a disk at the origin in $\mathbb{C}$ with coordinate t . Let $\eta : \mathcal{E} \rightarrow \Delta_t$ be a flat family of curves of arithmetic genus 1 satisfying:

- (i.) $\eta^{-1}(0) \cong C_j$.
- (ii.) $\eta^{-1}(t \neq 0)$ is irreducible, reduced, and (at worst) nodal.

For each $1 \leq i \leq d$, let $\mathcal{G}_i = \mathcal{H}_i \subset \mathcal{E}$ be the open subset of $\mathcal{E}$ on which the morphism η is *smooth*. Consider the fiber product:

$$X = \mathcal{G}_1 \times_{\Delta_t} \cdots \times_{\Delta_t} \mathcal{G}_d \times_{\Delta_t} \mathcal{H}_1 \times_{\Delta_t} \cdots \times_{\Delta_t} \mathcal{H}_d.$$

X is a nonsingular open set of the $2d$ -fold fiber product of $\mathcal{E}$ over Δ_t . Let $Y \subset X$ be the subset of points $y = (g_1, \dots, g_d, h_1, \dots, h_d)$ where the two divisors $\sum g_i$ and $\sum h_i$ are linearly equivalent on the curve $\eta^{-1}(\eta(y))$. Y is a nonsingular divisor in X .

Let $p \in C_j = \eta^{-1}(0)$ be a nonsingular point of C_j . Certainly $p \in \mathcal{G}_i, \mathcal{H}_i$ for all i . Let $\gamma : \Delta_t \rightarrow \mathcal{E}$ be any local holomorphic section of η such that $\gamma(0) = p$. Let V be a local holomorphic field of vertical tangent vectors to $\mathcal{E}$ on an open set containing p . The section γ and the vertical vector field V together determine local holomorphic coordinates (t, v) on $\mathcal{E}$ at p . Let $\phi_V : \mathcal{E} \times \mathbb{C} \rightarrow \mathcal{E}$ be the holomorphic flow of V defined locally near $(p, 0) \in \mathcal{E} \times \mathbb{C}$. The coordinate map

$$\psi : (t, v) \rightarrow \mathcal{E}$$

is determined by $\psi(t, v) = \phi_V(\gamma(t), v)$.

Local coordinates on X near the point $x_p = (p, \dots, p, p, \dots, p) \in X$ are given by

$$(t, v_1, \dots, v_d, w_1, \dots, w_d).$$

The coordinate map is determined by:

$$\psi_X(t, v_1, \dots, v_d, w_1, \dots, w_d) = (\psi(t, v_1), \dots, \psi(t, v_d), \psi(t, w_1), \dots, \psi(t, w_d)) \in X.$$

Note $x_p \in Y$. Let $f(t, v_1, \dots, v_d, w_1, \dots, w_d)$ be a local equation of Y at x_p . Since f is identically 0 on the line $(t, 0, \dots, 0, 0, \dots, 0)$,

$$(1) \quad \forall k \geq 0, \quad \frac{\partial^k f}{\partial t^k} \Big|_{x_p} = 0.$$

The tangent directions in the plane $t = 0$ correspond to divisors on the fixed curve C_j . Here, it is well known (up to $\mathbb{C}^*$ -factor)

$$(2) \quad \frac{\partial f}{\partial v_i} \Big|_{x_p} = +1, \quad \frac{\partial f}{\partial w_i} \Big|_{x_p} = -1.$$

Equations (1) and (2) are the only properties of f that will be used.

Let $\hat{\eta} : \hat{\mathcal{E}} \rightarrow \Delta_t$ be the family obtained by blowing-up $\mathcal{E}$ at p and adding $3d - 1$ -marking. Let $\mu : \hat{\mathcal{E}} \rightarrow \mathbf{P}^2$ be a morphism. Let $\hat{\eta}^{-1}(0) = D = C_j \cup \mathbf{P}^1$. Assume the following conditions are satisfied:

- (i.) $\mu, \hat{\eta}$, and the $3d - 1$ markings determine a family of Kontsevich stable pointed maps to $\mathbf{P}^2$.

- (ii.) The markings of D are distributed according to Φ .
- (iii.) $\deg(\mu|_{C_j}) = 0$, $\deg(\mu|_{\mathbf{P}^1}) = d$.

Let L_1, L_2 be general divisors of $\mu^*(\mathcal{O}_{\mathbf{P}^2}(1))$ that each intersect $\mathbf{P}^1$ transversely at d distinct points. For $1 \leq \alpha \leq 2$, L_α breaks into holomorphic sections $s_{\alpha,1} + \dots + s_{\alpha,d}$ of $\hat{\eta}$ over a holomorphic disk at $0 \in \Delta_t$. These sections $s_{\alpha,i}$ ($1 \leq \alpha \leq 2$, $1 \leq i \leq d$) determine a map $\lambda : \Delta_t \rightarrow Y$ locally at $0 \in \Delta_t$. Let an affine coordinate on $\mathbf{P}^1$ be given by ξ corresponding to the normal direction

$$(3) \quad \frac{d\gamma}{dt}|_{t=0} + \xi \cdot V(p).$$

Let $s_{1,i}(0) = \nu_i \in \mathbb{C} \subset \mathbf{P}^1$, $s_{2,i}(0) = \omega_i \in \mathbb{C} \subset \mathbf{P}^1$ be given in terms of the affine coordinate ξ . The map λ has the form

$$\lambda(t) = (t, \nu_1 t, \dots, \nu_d t, \omega_1 t, \dots, \omega_d t)$$

to first order in t (written in the coordinates determined by ψ_X). Equations (1), (2), and the condition $f(\lambda(t)) = 0$ implies

$$(4) \quad \sum_{i=1}^d \nu_i = \sum_{i=1}^d \omega_i.$$

$L_1 \cap \mathbf{P}^1$ is a degree d polynomial with roots at ν_i . Condition (4) implies that the sums of the roots (in the coordinates (3)) of general elements of the linear series $\mu|_{\mathbf{P}^1}$ are the same. Therefore, a constant K exists with the following property. If

$$\beta_0 + \beta_1 \xi + \dots + \beta_{d-1} \xi^{d-1} + \beta_d \xi^d$$

is an element of the linear series $\mu|_{\mathbf{P}^1}$, then $\beta_{d-1} + K \cdot \beta_d = 0$. The vanishing sequence at $\xi = \infty$ is therefore $\{0, \geq 2, *\}$. The point $\xi = \infty$ is the intersection $C_j \cap \mathbf{P}^1$.

Suppose $\tilde{\eta} : \tilde{\mathcal{E}} \rightarrow \Delta_t$ is obtained from $\mathcal{E}$ by a sequence of n blow-ups over p . The fiber $\tilde{\eta}^{-1}(0)$ is assumed to be C_j union a chain of $\mathbf{P}^1$'s of length n . Each blow-up occurs in the exceptional divisor of the previous blow-up. Let $\mathbf{P}$ denote the extreme exceptional divisor. Let $\mu : \tilde{\mathcal{E}} \rightarrow \mathbf{P}^2$ be of degree d on $\mathbf{P}$ and degree 0 on the other components of the special fiber $\tilde{\eta}^{-1}(0)$. Let there be $3d - 1$ markings as before. It must be again concluded that the linear series on $\mathbf{P}$ has vanishing sequence $\{0, \geq 2, *\}$ at the node.

Let γ be section of η such that the lift of γ to $\tilde{\eta}$ meets P . Let the coordinates (t, v) on $\mathcal{E}$ be determined by this γ (and any V). An affine coordinate ξ is obtained on $\mathbf{P}$ in the follow manner. Let γ_ξ be the section of η determined in (t, v) coordinates by

$$\gamma_\xi(t) = (t, \xi t^n).$$

Let $\tilde{\gamma}_\xi$ be the lift of γ_ξ to a section of $\tilde{\eta}$. The association

$$\mathbb{C} \ni \xi \mapsto \tilde{\eta}(0) \in \mathbf{P}$$

is an affine coordinate on $\mathbf{P}$. Let L_1, L_2 be divisors in the linear series μ intersecting $\mathbf{P}$ transversely. As before, L_α breaks into holomorphic sections $s_{\alpha,1}$. Let $s_{1,i} = \nu_i \in \mathbb{C} \subset \mathbf{P}$, $s_{2,i} = \omega_i \in \mathbb{C} \subset \mathbf{P}$. As before, a map $\lambda : \Delta_t \rightarrow Y$ is obtained from the sections $s_{\alpha,i}$. In the coordinates determined by ψ_X ,

$$\lambda(t) = (t, \nu_1 t^n + O(t^{n+1}), \dots, \nu_d t^n + O(t^{n+1}), \omega_1 t^n + O(t^{n+1}), \dots, \omega_d t^n + O(t^{n+1})).$$

As before $f(\lambda(t)) = 0$. The term of leading order in t of $f(\lambda(t))$ is

$$\left(\sum_{i=1}^d \nu_i - \sum_{i=1}^d \omega_i \right) \cdot t^n.$$

This follows from equations (1) and (2). The vanishing sequence $\{0, \geq 2, *\}$ is obtained as before.

By definition, an element $[\mu] \in I(\Phi, j)$ can be obtained as the special fiber of family of Kontsevich stable maps where the domain is a smoothing of the node p . After resolving the singularity in the total space at the node p by blowing-up, a family $\tilde{\mathcal{E}}$ is obtained. The above results show the linear series on $\mathbf{P}^1$ has vanishing sequence $\{0, \geq 2, *\}$ at p . $\square$

The markings play no role in the preceding proof. An identical argument establishes the following:

Lemma 2. *Let Φ be a stable, marked, weighted tree with distinguished vertex satisfying $e = 0$ and $w_i = d$ for some i . Let k be the number of non-distinguished vertices of Φ . Let $j \in \overline{M}_{1,1}$. Let $I(\Phi, j) = \overline{W}_1(d) \cap U_j(\Phi)$. The dimension of $I(\Phi, j)$ is bounded by $\dim I(\Phi, j) \leq 6d - k - 2$*

Lemma 3. *Let Ω be a stable, marked, weighted graph with 1 circuit. Let v_i be a non-circuit vertex with weight $w_i = d$ (this implies $e = 0$). Let $k + 1$ be the total number of vertices of Ω . Let $I(\Omega, \infty) = \overline{W}_1(d) \cap U_\infty(\Omega)$. The dimension of $I(\Omega, \infty)$ is bounded by $\dim I(\Omega, \infty) \leq 6d - k - 2$.*

The vanishing sequence $\{0, \geq 2, *\}$ condition reduces the dimensions of $U_C(\Phi)$, $U_\infty(\Omega)$ by 2.

4. THE NUMBERS $E_{d,j}$

The space of maps $\overline{M}_1(d)$ is equipped with $3d - 1$ evaluation maps corresponding to the marked points. For $i \in S_d$, let $e_i : \overline{W}_1(d) \rightarrow \mathbf{P}^2$ be the restriction of the i^{th} evaluation map to $\overline{W}_1(d)$. let $\mathcal{L}_i = e_i^*(\mathcal{O}_{\mathbf{P}^2})$. Let

$$Z = c_1(\mathcal{L}_1)^2 \cap \dots \cap c_1(\mathcal{L}_{3d-1})^2$$

Let $\pi_{\overline{W}} : \overline{W}_1(d) \rightarrow \overline{M}_{1,1} \cong \mathbf{P}^1$ be the restriction of π to $\overline{W}_1(d)$. Let

$$T = c_1(\pi_{\overline{W}}^*(\mathcal{O}_{\mathbf{P}^1}(1))).$$

Note $\overline{W}_1(d)$ is an irreducible, projective scheme of dimension $6d - 1$. The top intersection of line bundles on $\overline{W}_1(d)$, $Z \cap T$, is an integer.

Lemma 4.

$$\begin{aligned} \forall j \neq 0, 1728, \infty, \quad & Z \cap T = E_{d,j} , \\ j = 0, \quad & Z \cap T = 3 \cdot E_{d,0} , \\ j = 1728, \quad & Z \cap T = 2 \cdot E_{d,1728} . \end{aligned}$$

Proof. Via pull-back, lines in $\mathbf{P}^2$ yield representative classes of $c_1(\mathcal{L}_i)$. Therefore $3d - 1$ general points in $\mathbf{P}^2$, $\overline{x} = (x_1, \dots, x_{3d-1})$, determine a representative cycle $Z_{\overline{x}}$ of the the class Z . Let $\infty > j \in \overline{M}_{1,1}$. Let π_W be the restriction of π to $W_1(d)$. It is first established for a general representative $Z_{\overline{x}}$,

$$(5) \quad Z_{\overline{x}} \cap \pi_W^{-1}(j) \subset \pi_W^{-1}(j).$$

The statement (5) is proven by considering the quasi-projective strata of $\overline{M}_{C_j}(d)$.

Note $\pi_W^{-1}(j)$ is the strata $U_{C_j}(\Gamma, c, \overline{w})$ where $(\Gamma, c, \overline{w})$ is the trivial, stable, marked, weighted tree with distinguished vertex. Assume now $(\Gamma, c, \overline{w})$ is not the trivial tree. By the equations for the dimension of $(\Gamma, c, \overline{w})$

$$\dim U_{C_j}(\Gamma, c, \overline{w}) \leq 6d - 3$$

unless $e = 0$ and $k = 1, 2$. Since the linear series determined by the evaluation maps are base point free, the general intersection (5) will miss all loci of dimension less than $6d - 2$.

It remains to consider the trees $(\Gamma, c, \overline{w})$ where $e = 0$ and $k = 1, 2$. If $k = 1$, $(\Gamma, c, \overline{w}) = \Phi$ satisfies the conditions of Lemma (1). By Lemma (1),

$$\dim I(\Phi, j) \leq 6d - 3.$$

Hence, the general intersection (5) will miss all the loci $U_C(\Phi, c, (0, d))$.

If $k = 2$, there are two cases to consider. If there exists a vertex of weight d , then $(\Gamma, c, \overline{w}) = \Phi$ satisfies the conditions of Lemma (2). By Lemma (2),

$$\dim I(\Phi, j) \leq 6d - 4.$$

If $w_1 + w_2 = d$ is a positive partition, then the image of every map $[\mu] \in U_C(\Gamma, c, \overline{w})$ is the union of two rational curves of degrees w_1 and w_2 . No such unions pass through $3d - 1$ general points. The proof of claim (5) is complete.

For $\infty > j \neq 0, 1728$, $\pi_W^{-1}(j)$ is a reduced, irreducible divisor of $W_1(d)$. Since the linear series determined by the evaluation maps are base point free, the general intersection cycle

$$(6) \quad Z_{\overline{x}} \cap \pi_W^{-1}(j)$$

is a reduced collection of $Z \cap T$ points. The general intersection cycle (6) also consists exactly of the reduced, nodal, degree d elliptic plane curves with j -invariant j passing through the points $\overline{x}$.

The argument for $j = 0, 1728$ is identical except that $\pi_W^{-1}(0)$, and $\pi_W^{-1}(1728)$ are divisors in $W_1(d)$ with multiplicity 3, 2 respectively. These multiplicities arise from the extra automorphisms for $j = 0, 1728$. Therefore the cycle (6) is a collection of

$$\begin{aligned} & \frac{1}{3} \cdot Z \cap T, \\ & \frac{1}{2} \cdot Z \cap T \end{aligned}$$

triple and double points respectively. $\square$

It remains to evaluate $Z \cap T$.

Lemma 5. $Z \cap T = \binom{d-1}{2} N_d$.

Proof. It is first established for a general representative $Z_{\overline{x}}$,

$$(7) \quad Z_{\overline{x}} \cap \pi_W^{-1}(\infty) \subset \pi_W^{-1}(\infty).$$

The statement (7) is proven by considering the quasi-projective strata of $\overline{M}_\infty(d)$.

By arguments of Lemma (4), all the loci $U_\infty(\Gamma, c, \overline{w})$ where $(\Gamma, c, \overline{w})$ is not the trivial tree are avoided in the general intersection (7). Only the strata $U_\infty(\Lambda, \overline{w})$ remain to be considered. Let $k + 1 \geq 2$ be the total number of vertices of Λ . By the equations for the dimensions of $U_\infty(\Lambda, \overline{w})$,

$$\dim U_\infty(\Lambda, \overline{w}) \leq 6d - 2 - k \leq 6d - 3$$

unless all the circuit vertices have weight zero. If all circuit vertices have weight zero, $k + 1 \geq 3$. Now

$$\dim U_\infty(\Lambda, \overline{w}) \leq 6d - k \leq 6d - 3$$

unless $k = 2$.

Only one stable, marked, weighted, graph with 1-circuit $(\Lambda, \overline{w})$ need be considered. Vertices c_1, c_2 form a weightless circuit. Vertex v_3 is connected to c_2 and $w_3 = d$. $(\Lambda, \overline{w}) = \Omega$ satisfies the conditions of Lemma (3). Therefore,

$$\dim I(\Omega, \infty) \leq 6d - 4.$$

Claim (7) is now proven.

The divisor $\pi_W^{-1}(\infty)$ is reduced and irreducible in $W_1(d)$. As above,

$$(8) \quad Z_{\overline{x}} \cap \pi_W^{-1}(\infty)$$

is a reduced collection of $Z \cap T$ points. The general intersection cycle (8) also consists exactly of degree d maps of the 1-nodal rational curve C_∞ passing through $\overline{x}$. The image of such a map must be one of the N_d degree d , nodal, rational plane curves passing through $\overline{x}$. The number of distinct birational maps (up to isomorphism) from C_∞ to a $\binom{d-1}{2}$ -nodal plane curve is exactly $\binom{d-1}{2}$. Therefore, $Z \cap T = \binom{d-1}{2} N_d$. $\square$

REFERENCES

- [Al] V. Alexeev, *Moduli Spaces $M_{g,n}(W)$ For Surfaces*, preprint 1994.
- [A] P. Aluffi, *How Many Smooth Plane Cubics With Given j -Invariant Are Tangent To 8 Lines In General Position?*, Contemporary Mathematics 123, 1991, p.15-29.
- [D-I] P. Di Francesco and C. Itzykson, *Quantum Intersection Rings*, preprint 1994.
- [K] M. Kontsevich, *Enumeration of Rational Curves Via Torus Actions*, preprint 1994.
- [K-M] M. Kontsevich and Y. Manin, *Gromov-Witten Classes, Quantum Cohomology, and Enumerative Geometry*, preprint 1994.
- [P] R. Pandharipande, *Notes On Kontsevich's Compactification Of The Moduli Space Of Maps*, Course Notes, 1995.

Department of Mathematics, University of Chicago

rahul@math.uchicago.edu